

Aytu Products Savings Offer



Present to your pharmacist to apply the savings offer

100% of eligible patients with commercial non-governmental insurance may pay as low as*



— or —



Per prescription if “covered” on insurance plan and where deductible has been met

Per prescription if “not covered” on insurance plan or if the deductible has not been met. No prior authorization required

*Terms and conditions apply.

Pharmacist Instructions:

1. Visit www.aytucoupon.com
2. Click “I’m a pharmacy”
3. Input necessary information and press submit
4. Utilize the offer to assist patient

The ACE Customer Excellence Team

If you experience any pharmacy-related issues, call our dedicated staff at 1-888-AYTURxC (298-8792). Support available Monday through Friday, 9:00 am to 7:30 pm ET (except holidays).

Retail Copay Offer Terms and Conditions: This program provides non-government copay assistance with out-of-pocket costs for eligible patients. By using this offer, patient and pharmacist understand and agree to comply with these terms and conditions. Offer may only be used by eligible residents of the U.S. at participating pharmacies and may not be redeemed at government-subsidized clinics. Offer user must be at least 18 years of age to use for themselves or on behalf of a minor. Patient age or insurance restrictions may apply.

Offer limited to one per person and is not transferrable. No substitutions are permitted. Offer eligible only with valid prescription, has no cash value, and cannot be combined with any free trial, discount, prescription savings card or other offer. This offer is not insurance. Valid only for patients with commercial insurance and NOT valid for prescriptions eligible to be reimbursed in whole or in part by Medicaid, Medicare (including Medicare Advantage and Part D plans), or any other federally or state funded healthcare benefit program, or by commercial plans or other health or pharmacy benefit programs that reimburse for entire cost of the prescription drug or prohibit offer’s use. Medicare Part D enrollees who are in the prescription drug coverage gap are not eligible for offer. Void where prohibited by law, taxed or restricted. It is illegal to sell, purchase, trade, or counterfeit, or offer to sell, purchase, trade or counterfeit the offer.

Patient, pharmacist and prescriber agree not to seek reimbursement for all or any part of the benefit received by patient through the offer. Certain information pertaining to use of the offer will be shared with Aytu BioPharma, the sponsor of the offer, and its affiliates. The information disclosed will include the patient copay ID, pharmacy demographics, prescriber information, and details relating to the claim, such as copay amount, insurance details, and therapy received. For more information, please see the Aytu BioPharma privacy policy.

Aytu BioPharma reserves the right to rescind, revoke or amend the offer at any time without notice.

Please see Important Safety Information on the following pages and [click here](#) for Medication Guide.

Most Important Information about ADZENYS XR-ODT

Abuse, misuse, and addiction. ADZENYS XR-ODT has a high chance for abuse and misuse and may lead to substance use problems, including addiction. Misuse and abuse of ADZENYS XR-ODT, other amphetamine containing medicines, and methylphenidate containing medicines, can lead to overdose and death. The risk of overdose and death is increased with higher doses of ADZENYS XR-ODT or when it is used in ways that are not approved, such as snorting or injection.

- ADZENYS XR-ODT may lead to physical dependence after prolonged use, even if taken as directed by your healthcare provider.
- Do not give ADZENYS XR-ODT to anyone else.
- Keep ADZENYS XR-ODT in a safe place and properly dispose of any unused medicine.

• **Risks for people with serious heart disease.** Sudden death has happened in people who have heart defects or other serious heart disease. Your healthcare provider should check you or your child carefully for heart problems before starting ADZENYS XR-ODT. Tell your healthcare provider if you or your child has any heart problems, heart disease, or heart defects.

Call your healthcare provider or go to the nearest hospital emergency room right away if you or your child has any signs of heart problems such as chest pain, shortness of breath, or fainting while taking ADZENYS XR-ODT.

- **Increased blood pressure and heart rate.** Your healthcare provider should check you or your child's blood pressure and heart rate regularly during treatment with ADZENYS XR-ODT.
- **Mental (Psychiatric) problems:** Tell your healthcare provider about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression. **Call your healthcare provider right away if you or your child have any new or worsening mental symptoms or problems while taking ADZENYS XR-ODT, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.**

What is ADZENYS XR-ODT: ADZENYS XR-ODT is a central nervous system (CNS) stimulant prescription medicine. **It is used for the treatment of Attention Deficit Hyperactivity Disorder (ADHD).** ADZENYS XR-ODT may help increase attention and decrease impulsiveness and hyperactivity in patients with ADHD. It is not known if ADZENYS XR-ODT is safe and effective in children under 6 years of age. **ADZENYS XR-ODT is a federally controlled substance (CII) because it contains amphetamine that can be a target for people who abuse prescription medicines or street drugs.** Keep ADZENYS XR-ODT in a safe place to prevent it from theft. Never give your ADZENYS XR-ODT to anyone else because it may harm them. Selling or giving away ADZENYS XR-ODT may harm others and is against the law.

Who should not take ADZENYS XR-ODT: ADZENYS XR-ODT should not be taken if you or your child:

- Are allergic to amphetamine, or any of the ingredients in ADZENYS XR-ODT
- Are taking or have taken within the past 14 days an anti-depression medicine called monoamine oxidase inhibitor or MAOI

What should I tell my healthcare provider before taking ADZENYS XR-ODT?

- All health conditions (or family history) of heart problems or high blood pressure; mental problems; circulation problems; kidney problems; if you, your child, or family member have or had repeated movements or sounds (tics) or Tourette's syndrome; if you or your child are pregnant or plan to become pregnant; or if you or your child is breastfeeding or plan to breastfeed.
- All medications that you or your child takes.
- Any anti-depression medicines including MAOIs
- Do not start any new medicine while taking ADZENYS XR-ODT without talking to your healthcare provider first.

Avoid drinking alcohol while taking ADZENYS XR-ODT.

The most common side effects of ADZENYS XR-ODT include decreased appetite and problems sleeping. ADZENYS XR-ODT can cause serious side effects, including:

- **Slowing growth (height and weight) in children.** Children should have their height and weight checked often during treatment with ADZENYS XR-ODT.
- **Circulation problems in fingers and toes.** [Peripheral vasculopathy, including Raynaud's phenomenon]: fingers and toes may feel numb, cool, painful and/or change from pale to blue or red. **Call your healthcare provider right away if you or your child have any unexplained wounds appearing on fingers or toes while taking ADZENYS XR-ODT.**
- **New or worsening tics or worsening Tourette's syndrome.** Tell your healthcare provider if you or your child get any new or worsening tics or worsening Tourette's syndrome during treatment with ADZENYS XR-ODT.
- Common side effects of ADZENYS XR-ODT **in children 6 to 12 years of age** also include: stomach pain, extreme mood change, vomiting, nervousness, nausea, and fever.
- Common side effects of ADZENYS XR-ODT **in children 13 to 17 years of age** also include: stomach pain and weight loss.
- Common side effects of ADZENYS XR-ODT **in adults** also include: dry mouth, headache, weight loss, nausea, anxiety, restlessness, dizziness, fast heartbeat, diarrhea, weakness, and urinary tract infections.

These are not all the possible side effects of ADZENYS XR-ODT.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Please click [here](#) for Full Medication Guide.

Most Important Information about COTEMPLA XR-ODT

Abuse, misuse, and addiction. COTEMPLA XR-ODT has a high chance for abuse and misuse and may lead to substance use problems, including addiction. Misuse and abuse of COTEMPLA XR-ODT, other methylphenidate containing medicines, and amphetamine containing medicines, can lead to overdose and death. The risk of overdose and death is increased with higher doses of COTEMPLA XR-ODT or when it is used in ways that are not approved, such as snorting or injection.

- **Risks for people with serious heart disease.** Sudden death has happened in people who have heart defects or other serious heart disease. Your healthcare provider should check your child carefully for heart problems before starting COTEMPLA XR-ODT. Tell your healthcare provider if your child has any heart problems, heart disease, or heart defects. **Call your healthcare provider or go to the nearest hospital emergency room right away if your child has any signs of heart problems such as chest pain, shortness of breath, or fainting while taking COTEMPLA XR-ODT.**
- **Increased blood pressure and heart rate.** Your healthcare provider should check your child's blood pressure and heart rate regularly during treatment with COTEMPLA XR-ODT.
- **Mental (Psychiatric) problems:** Tell your healthcare provider about any mental problems your child has, or about a family history of suicide, bipolar illness, depression, new psychotic symptoms (such as hearing voices, or seeing or believing things that are not real), or new manic symptoms. **Call your healthcare provider right away if your child has any new or worsening mental symptoms or problems while taking COTEMPLA XR-ODT, especially seeing or hearing things that are not real, believing things that are not real, or are suspicious.**

What is COTEMPLA XR-ODT: COTEMPLA XR-ODT is a central nervous system (CNS) stimulant prescription medicine used for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in children under 6 to 17 years of age. COTEMPLA XR-ODT may help increase attention and decrease impulsiveness and hyperactivity in children 6 to 17 years of age with ADHD. It is not known if COTEMPLA XR-ODT is safe and effective in children under 6 years of age.

You should avoid drinking alcohol during treatment with COTEMPLA XR-ODT.

COTEMPLA XR-ODT is a federally controlled substance (CII) because it contains methylphenidate that can be a target for people who abuse prescription medicines or street drugs. Keep COTEMPLA XR-ODT in a safe place to protect it from theft. Never give your COTEMPLA XR-ODT to anyone else because it may cause death or harm them. Selling or giving away COTEMPLA XR-ODT may harm others and is against the law.

Do not give COTEMPLA XR-ODT to your child if they are: COTEMPLA XR-ODT should not be taken if you or your child:

- Are allergic to methylphenidate or any of the ingredients in COTEMPLA XR-ODT
- Are taking or have taken within the past 14 days a medicine used to treat depression called a monoamine oxidase inhibitor MAOI

Before taking COTEMPLA XR-ODT tell your child's healthcare provider about all medical conditions, including if your child:

- Has heart problems, heart disease, heart defects, or high blood pressure
- Has mental problems including psychosis, mania, bipolar illness, or depression
- Has circulation problems in fingers or toes
- Have eye problems, including increased pressure in your eye, glaucoma, or problems with your close-up vision (farsightedness)
- Have, or had repeated movements or sounds (tics) or Tourette's syndrome, or have a family history of tics or Tourette's syndrome
- Is pregnant or plans to become pregnant. It is not known if COTEMPLA XR-ODT will harm the unborn baby.
- Is breastfeeding or plans to breastfeed. COTEMPLA XR-ODT passes into breast milk. You and your healthcare provider should decide if your child will take COTEMPLA XR-ODT or breastfeed.

Tell your healthcare provider about all of the medicines your child takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements. COTEMPLA XR-ODT and some medicines may interact with each other and cause serious side effects. Sometimes the doses of other medicines will need to be adjusted during treatment with COTEMPLA XR-ODT. Your healthcare provider will decide whether COTEMPLA XR-ODT can be taken with other medicines. **Especially tell your healthcare provider if your child takes anti-depression medicines including MAOIs.**

 **Cotempla XR-ODT** 
(methylphenidate) Extended-Release Orally Disintegrating Tablets
8.6 mg, 17.3 mg, 25.9 mg

Most Important Information about COTEMPLA XR-ODT (cont'd)

Do not start any new medicine during treatment with COTEMPLA XR-ODT without talking to your healthcare provider first.

The most common side effects of COTEMPLA XR-ODT include decreased appetite and problems sleeping. COTEMPLA XR-ODT can cause serious side effects, including:

- **Painful or prolonged erections (priapism).** If your child develops priapism, get medical help right away.
- **Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon):** fingers and toes may feel numb, cool, painful and/or change from pale to blue or red. **Tell your healthcare provider if your child has numbness, pain, skin color change, sensitivity to temperature in their fingers or toes, or any signs of unexplained wounds appearing on fingers or toes during treatment with COTEMPLA XR-ODT.**
- **Slowing of growth (height and weight in children).** Children should have their height and weight checked often during treatment with COTEMPLA XR-ODT. COTEMPLA XR-ODT treatment may be stopped if your child is not gaining weight or height.
- **Eye problems (increased pressure in the eye and glaucoma).** Call your healthcare provider right away if your child develops changes in their vision or eye pain, swelling, or redness.
- **New or worsening tics or worsening Tourette's syndrome.** Tell your healthcare provider if your child gets any new or worsening tics or worsening Tourette's syndrome during treatment with COTEMPLA XR-ODT.
- The most common side effects of methylphenidate products include: decreased appetite, trouble sleeping, nausea, vomiting, indigestion, stomach pain, weight loss, anxiety, dizziness, irritability, mood swings, increased heart rate, increased blood pressure.

These are not all the possible side effects of COTEMPLA XR-ODT.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Please [click here](#) for Full Medication Guide.


Cotempla XR-ODT 
(methylphenidate) Extended-Release Orally Disintegrating Tablets
8.6 mg, 17.3 mg, 25.9 mg